

## References

*The learning package has been developed through a partnership consisting of the Workforce Development Branch of the Department of Health, Health Networks Branch sponsors of the WA Smoke Free Health Working Party and the National Drug Research Institute, Curtin University of Technology. It has also drawn on learning packages developed by a group consisting of the above collaborators, the WA Country Health Services and the WA Drug and Alcohol Office.*

- Addy, D., Skinner, N., Shoobridge, J., Freeman, T., Roche, A.M., Pidd, K., Watts, S., (2004). *The Work Practice Questionnaire: A training evaluation measurement tool for the alcohol and other drugs field*. Adelaide: National Centre for Education and Training on Addiction
- Australian Institute of Health and Welfare. (2008). *2007 National Drug Strategy Household Survey: First Results*. Canberra: Australian Institute of Health and Welfare.
- Babor, T. F. (1990). Brief intervention strategies for harmful drinkers: New directions for medical education. *Canadian Medical Association Journal*, 143(10), 1070-1076.
- Balfour, D. J. & Fagerstrom, K. O. *Pharmacol Ther* 1996 72:51-81
- Bjartveit, K., & Tverdal, A. (2005). Health consequences of smoking 1-4 cigarettes per day. *Tobacco Control*, 14(5), 315-320.
- Carter, S., Borland, R. & Chapman, S. (2001) *Finding the strength to kill your best friend: Smokers talk about smoking and quitting*. Sydney: Australian Smoking Cessation Consortium and GlaxoSmithKline Consumer Healthcare
- Coase, P., & Miller, J. (2007). *ASSAD Smoking Report 2005*: Prepared for Drug and Alcohol Office WA. Perth: TNS Social Research Report No. 70409
- Collins, D. J., Lapsley, H. M., & University of New South Wales. (2008). *The costs of tobacco, alcohol and illicit drug abuse to Australian society in 2004/2005*. Canberra: Commonwealth of Australia
- Dawe, S., Loxton, N., Hides, L., Kavanagh, D. & Mattick, R. P. (2002). *Diagnostic and screening instruments for alcohol and other drugs and other psychiatric disorders*. Revised National Drug Strategy Monograph. Canberra: Commonwealth Department of Human Services and Health.
- Department of Health. (2008). Clinical guidelines and procedures for the management of nicotine dependent inpatients. Perth: Department of Health.
- Department of Health Western Australia. (2004) *The Quit Book - A Guide to Quitting Smoking, Quit WA*. Perth: Department of Health
- Doll, R., Peto, R., Boreham, J., & Sutherland, I. (2004). Mortality in relation to smoking: 50 years' observations on male British doctors. *British Medical Journal*, 328(7455), 1519.
- Emmons, K., & Goldstein, M. (1992). Smokers who are hospitalized: A window of opportunity for cessation interventions. *Preventative Medicine*, 21 (2), 262-269.
- Fagerstrom, K. O. (1978). Measuring degree of physical dependence to tobacco smoking with reference to individualization of treatment. *Addictive Behaviors*. 3, 34, 235-241
- Fagerstrom, K. O., & Schneider, N. G. (1989). Measuring nicotine dependence: a review of the Fagerstrom Tolerance Questionnaire. *Journal of Behavioral Medicine*, 12(2):159-182.
- Fiore, M. C., Bailey, W. C., Cohen, S. J., Dorfman, S. F., Goldstein, M. G., Gritz, E. R., Heyman, R. B., Holbrook, J., Jaen, C. R., Kottke, T. E., Lando, H. A., Mecklenburg, R. E., Mullen, P. D., Nett, L. M., Robinson, L., Stitzer, M. L., Tommasello, A. C., Villejo, L., Wewers, M. E. (2000). *Treating Tobacco Use and Dependence: Clinical Practice Guidelines*. Rockville, MD: U.S. Department of Health and Human Services, Public Health Service.

- Fong, G. T., Hammond, D., Laux, F. L., Zanna, M. P., Cummings, K. M., Borland, R., & Ross H. (2004). The near-universal experience of regret among smokers in four countries: Findings from the International Tobacco Control Policy Evaluation Survey. *Nicotine & Tobacco Research* 6(3), 341 - 351.
- Higgins-Biddle, J., & Babor, T. (2006) "Reducing risky drinking via screening and brief intervention," *Alcoholism & Drug Abuse Weekly*, vol. 8 (37), 5.
- Hill, D., & Borland, R. (1991). Adults' accounts of onset of regular smoking: influences of school, work and other settings. *Public Health Reports*, 106(2), 181-185.
- Hutchison, D., Shepstone, L., Moots, R., Lear, J., & Lynch, M. (2001). Heavy cigarette smoking is strongly associated with rheumatoid arthritis (RA), particularly in patients without a family history of RA. *Annals of Rheumatic Diseases*; 60: 223-227
- Kaner, E. F. S., Beyer, F., Dickinson, H. O., Pienaar, E., Campbell, F., Schlesinger, C., Heather, N., Saunders, J., & Burnand, B. (2007). Effectiveness of brief alcohol interventions in primary care populations. *Cochrane Database of Systematic Reviews*, Issue 2. Art No.: CD004148. DOI: 10.1002/14651858.CD004148.pub3.
- Kelly, S., Thornton, J., & Lyratzopolous, G, et al: Smoking and blindness, *British Medical Journal* 2004; 328: 537- 8
- Lancaster, T., Steed, L., Silagy, C., & Sowden, A. (2000). Effectiveness of interventions to help people stop smoking: *Findings from the Cochrane Library*. 321, 355-358.
- Mahonen, M. S., McElduff, P., Dobson, A. J., Kuulasmaa, K. A., Evans, A. E., & Project, W. M. (2004). Current smoking and the risk of non-fatal myocardial infarction in the WHO MONICA Project populations. *Tobacco Control*, 13(3), 244-250.
- McCambridge, J., & Day, M. (2008). Randomized controlled trial of the effects of completing the Alcohol Use Disorders Identification Test questionnaire on self-reported hazardous drinking. *Addiction*, 103, 241-248.
- Miller, W.R., & Hester, R. K. (1986). The effectiveness of alcoholism treatment: what research reveals. In W.R. Miller, and N. Heather, (eds). *Treating addictive behaviours: process of change*. New York: Plenum Press.
- Miller, W. R., & Rollnick, S. (1991). *Motivational interviewing: Preparing people to change addictive behaviour*. New York: Guildford Press.
- Miller, W. R., and Rollnick, S. (2002). *Motivational interviewing*. (2nd ed.). New York: Guildford Press.
- Miller, M., & Wood, L. (2001). *Smoking cessation interventions: Review of evidence and implications for best practice in health care settings*. Canberra: Australian Government, Department of Health and Ageing.
- Mosher, V., Davis, J., Mulligan, D., & Iber, F.L. (1975). Comparison of outcome in a 9-day and 30-day alcoholism treatment program. *Journal of Studies on Alcohol*, 36, 1277 - 1281.
- National Health and Medical Research Council. (1997). *The health effects of passive smoking: A scientific information paper*. Canberra: NHMRC.
- Owen, N. (1985). Advising patients to stop smoking. *Medical Journal of Australia*, 142, 176-178.
- Page, R.D., and Shaub, L.H. (1979). Efficacy of a three versus a five week alcohol treatment program. *International Journal of the Addictions*, 14, 697-714
- Peters, M. J., Morgan, L. C., & Gulch, L. (2004). Smoking cessation and elective surgery: The cleanest cut. *Medical Journal of Australia*, 180, 317-318.
- Peto, R. (1994). Smoking and death: the past 40 years and the next 40. *British Medical Journal*, 309(6959), 937-939.

- Resnicow, K., Dilorio, C., Soet, J. E., Borrelli, B., Ernst, D., and Hecht, J. (2002). Motivational Interviewing in Health Promotion: It sounds like something is changing. *Health Psychology*, 21, 444-451.
- Rice, V. H., & Stead, L. F. (1999). Nursing interventions for smoking cessation. *Cochrane Database of Systematic Reviews*, Issue 3. Art. No.: CD001188. DOI: 10.1002/14651858.CD001188.pub3
- Rigotti, N. A., Munafo, M. R., & Stead, L. F. (2007). Interventions for smoking cessation in hospitalised patients. *Cochrane Database of Systematic Reviews*, 3, CD001837.
- Rollnick, S., Mason, P., & Butler, C. (1999). *Health Behavior Change: A guide for Practitioners*, Churchill Livingstone, London.
- Royal College of Physicians (1992). *Smoking and the young*. London: Royal College of Physicians.
- Russell, M.A.H., Wilson, C., Taylor, C. and Baker, C. (1979). Effect of General Practitioners' advice against smoking. *British Medical Journal*, 2, 231-235.
- Stead, L. F., Bergson, G., & Lancaster, T. (2008). Physician advice for smoking cessation. *Cochrane Database of Systematic Reviews*, 2, 1-42.
- Stead LF, Perera R, Bullen C, Mant D, Lancaster T (2008) Nicotine replacement therapy for smoking cessation. In *Cochrane Database of Systematic Reviews: Reviews 2008*. John Wiley & Sons, Ltd Chichester, UK.
- Unwin, E., Codde, J., & Bartu, A. (2003). *The impact of tobacco smoking on the health of Western Australians*. Perth: Drug and Alcohol Office and the Epidemiology Branch, Department of Health, Western Australia.
- US Department of Health and Human Services (2001). *Smoking and women's health. A report of the Surgeon General*. Rockville: USDHHS.
- U.S. Department of Health and Human Services (2004). *The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health.
- Viegi, G., Scognamilio, A., Baldacci, S., Pistelli, F., & Carozzi, L. (2001). Epidemiology of Chronic Obstructive Pulmonary Disease (COPD). *Respiration*, 68, 4-19.
- White, V., & Scollo, M. (2003). How many children take up smoking each year in Australia? *Australian and New Zealand Journal of Public Health*, 27(3), 359-360.
- Wolfenden, L., Campbell, E., Walsh, R., & Wiggers, J. (2003). Smoking cessation interventions for in-patients: A selective review with recommendations for hospital-based health professionals. *Drug and Alcohol Review*, 22(4), 437-452.
- Zwar, N., Bell, J., Peters, M., Christie, M. & Mendelsohn, C. (2006). Nicotine and nicotine replacement therapy - the facts. *Pharmacist*, 25 (12), 969-973.