

References

- Australian Bureau of Statistics (ABS) (2015). Australian Demographic Statistics, Dec 2015. Retrieved from:
<http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/3101.0Dec%202015?OpenDocument>
- Bagnardi, V., M. Rota, E. Botteri, I. Tramacere, F. Islami, V. Fedirko, L. Scotti, et al (2015). *Alcohol Consumption and Site-Specific Cancer Risk: A Comprehensive Dose-Response Meta-Analysis*. Br J Cancer 112, no. 3: 580-93.
- English, D. R., Holman, C. D. J., Milne, E., Winter, M. G., Hulse, G. K., Codde, J. P., Bower, C. I., Corti, B., de Klerk, N., Knuiiman, M. W., Kurinczuk, J. J., Lewin, G. F., & Ryan, G. A. (1995). *The Quantification of Drug Caused Morbidity and Mortality in Australia, 1995 Edition* (Vol. 1). Canberra: Australian Government Publishing Service.
- Matthews, S., T. Chikritzhs, P. Catalano, T. Stockwell, and S. Donath (2002). *Trends in Alcohol-Related Violence in Australia, 1991/92-1999/00. National Alcohol Indicators, Bulletin No.5*. Perth: National Drug Research Institute, Curtin University of Technology, 2002.
- Ridolfo, B. and C. Stevenson (2001), *The quantification of drug-caused mortality and morbidity in Australia, 1998*. AIHW cat. no. PHE 29. 2001, Australian Institute of Health and Welfare: Canberra.
- Sherk A., Stockwell T., Rehm J., Dorocicz J., Shield K.D. (2017). *The International Model of Alcohol Harms and Policies (InterMAHP) v1.0: A comprehensive guide to the estimation of alcohol-attributable morbidity and mortality*. Victoria, BC: Centre for Addictions Research of British Columbia, University of Victoria
- World Health Organization (WHO). (2000). *International guide for monitoring alcohol consumption and related harm*. (WHO/MSD/MSB/00.4). Geneva: WHO Department of Mental Health and Substance Dependence, Noncommunicable Diseases and Mental Health Cluster, World Health Organization.